

uc3m

Universidad **Carlos III** de Madrid

Departamento de Matemáticas

Complex variable and transforms. Problems

Chapter 1: Complex variable

Section 1.5: Complex integration

Professors:

Domingo Pestana Galván

José Manuel Rodríguez García



1.5. COMPLEX INTEGRATION

5.1. Compute the following integrals:

- a) $\int_{\gamma} x dz$, where γ is the segment joining 0 with $1 + i$.
- b) $\int_{\gamma} x dz$, where γ is the circumference with radius r centered at 0.
- c) $\int_{\gamma} \frac{dz}{z}$, with $\gamma(t) = \cos t + 2i \sin t$, $t \in [0, 2\pi]$.
- d) $\int_{\gamma} \frac{dz}{z^2}$, with $\gamma(t) = \cos t + 2i \sin t$, $t \in [0, 2\pi]$.
- e) $\int_{\gamma} \frac{dz}{z^2 - 1}$, with $\gamma(t) = 1 + e^{it}$, $t \in [0, 2\pi]$.
- f) $\int_{\gamma} |z| \bar{z} dz$, where γ is the boundary of the half-disk $\{z \in \mathbf{C} : |z| \leq 1, \text{Im } z \geq 0\}$.
- g) $\int_{\gamma} |z - 1| |dz|$, where γ is the unit circumference (the boundary of the disk with radius 1 centered at 0).
- h) $\int_{\gamma} |z|^2 dz$, where γ is the square with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$ and $(0, 1)$.
- i) $\int_{\gamma} \frac{|z|}{|1 - z|^2} |dz|$, where γ is the circumference with radius r ($0 < r < 1$) centered at 0.
- j) $\int_{\gamma} \frac{dz}{|z - a|^2}$, where γ is the circumference with radius r centered at 0, and $a \in \mathbf{C}$ with $|a| \neq r$.

Hints: i) prove first that if $0 \leq r < R$, then

$$\frac{1}{R^2 - 2rR \cos t + r^2} = \frac{1}{R^2 - r^2} \left(1 + 2 \sum_{n=1}^{\infty} \left(\frac{r}{R} \right)^n \cos nt \right).$$

j) Recall that

$$|z - a|^2 = (z - a)(\bar{z} - \bar{a}) = (z - a)(r^2/z - \bar{a}).$$

5.2. Compute $\int_{\gamma} \bar{z} dz$, if γ is:

- a) $\gamma(t) = t^2 + it$, with $0 \leq t \leq 2$,
- b) the polygonal joining the points 0 with $2i$, and $2i$ with $4 + 2i$.

Can an holomorphic function exist with derivative \bar{z} ?

5.3. Compute $\int_{\gamma} \frac{dz}{\bar{z}^2}$, if γ is:

- a) $\gamma(t) = e^{i(\pi-t)}$, with $0 \leq t \leq \pi$,
- b) $\gamma(t) = e^{it}$, with $\pi \leq t \leq 2\pi$.

Do the answers of a) and b) imply that there exists an holomorphic function f with $f'(z) = 1/\bar{z}^2$?

5.4. a) Compute for each $n \in \mathbf{Z}$ and $a \in \mathbf{C}$ the integral:

$$I = \int_{\gamma} (z - a)^n dz,$$

where γ is any circumference with $a \notin \gamma$.

- b) Prove that there is no holomorphic function f on $\mathbf{C} \setminus \{0\}$ such that $f'(z) = 1/z$.
 c) If $\gamma = \{z \in \mathbf{C} : |z| = 2\}$, compute the following integral (by using the index):

$$\int_{\gamma} \frac{dz}{z^2 - 1}.$$

- d) If $\gamma = \{z \in \mathbf{C} : |z| = 3\}$, compute the following integral:

$$\int_{\gamma} \frac{2z^2 - 15z + 30}{z^3 - 10z^2 + 32z - 32} dz.$$

5.5. a) If $f : [a, b] \rightarrow \mathbf{C}$ is an integrable function, prove that:

$$\left| \int_a^b f(t) dt \right| \leq \int_a^b |f(t)| dt$$

Hint: If $\int_a^b f(t) dt = R e^{i\alpha}$, then $R = \int_a^b \operatorname{Re}(e^{-i\alpha} f(t)) dt$.

b) Let $D \subset \mathbf{C}$ be an open set, $f : D \rightarrow \mathbf{C}$ a continuous function and γ a curve contained on D . Prove that:

$$\left| \int_{\gamma} f(z) dz \right| \leq \int_{\gamma} |f(z)| |dz|.$$

- c) With the hypotheses in the previous item, prove that

$$\left| \int_{\gamma} f(z) dz \right| \leq M \operatorname{length} \gamma,$$

if $|f(z)| \leq M$ on γ .

- d) Prove that if $\gamma(t) = e^{it}$ with $t \in [0, \pi]$, we have

$$\left| \int_{\gamma} \frac{e^z}{z} dz \right| \leq \pi e.$$

- e) Prove that if γ is the unit circumference positively oriented, we have

$$\left| \int_{\gamma} \frac{\sin z}{z^2} dz \right| \leq 2\pi e.$$

It would have the same inequality if γ were oriented in the opposite direction?

5.6. a) If $f_n \rightarrow f$ uniformly on $\gamma([a, b])$, prove that $\int_{\gamma} f_n \rightarrow \int_{\gamma} f$ as $n \rightarrow \infty$.

Hint: Use item c) in the previous exercise.

b) Consider the curves $\gamma : [0, 2\pi] \rightarrow \mathbf{C}$ with $\gamma(t) = r e^{it}$ and $\gamma_n : [0, 2\pi] \rightarrow \mathbf{C}$ with $\gamma_n(t) = (1 - 1/n) r e^{it}$. If f is a continuous function on the closed disk centered at 0 with radius r , prove that:

$$\int_{\gamma} f(z) dz = \lim_{n \rightarrow \infty} \int_{\gamma_n} f(z) dz.$$

Hint: You can use item a).