

**uc3m**

Universidad **Carlos III** de Madrid

Departamento de Matemáticas

## Complex variable and transforms. Problems

Chapter 1: Complex variable

Section 1.6: Cauchy's integral formula

Professors:

Domingo Pestana Galván

José Manuel Rodríguez García



## 1.6. CAUCHY'S INTEGRAL FORMULA

**6.1.** Compute the following integrals:

$$\begin{aligned}
 & a) \int_{|z|=1} \frac{\cos z}{z} dz, \quad b) \int_{|z|=1} \frac{\sin z}{z^2} dz, \quad c) \int_{|z|=3} \frac{e^z + z}{z-2} dz, \\
 & d) \int_{|z|=2} \frac{z^2}{z-1} dz, \quad e) \int_{|z|=2} \frac{z^2-1}{z^2+1} dz, \quad f) \int_{|z|=2} \frac{dz}{z^2+2z-3}, \\
 & g) \int_{|z|=2} \frac{|z|e^z}{z^2} dz, \quad h) \int_{|z-1|=2} \frac{dz}{z^2-2i}, \quad i) \int_{|z|=2} \frac{dz}{z^2(z^2+16)}, \\
 & j) \int_{|z|=3/2} \frac{\sinh 5z}{(1+z^2)z^2} dz, \quad k) \int_{|z-z_0|=r} \frac{2z - \sin 2z + 2(z-z_0)\cos^2 z}{(z-z_0)^2} dz, \\
 & \quad l) \int_{|z-2|=1} \frac{e^z}{z} dz, \quad m) \int_{|z|=1} \frac{e^z}{z} dz, \quad n) \int_{|z|=1} \frac{\sin z}{z} dz, \\
 & o) \int_{|z|=1} \frac{e^{3z}}{(z-1/2)^5} dz, \quad p) \int_{|z|=2} \frac{z^2}{z^3-1} dz, \quad q) \int_{|z|=2} \frac{z^2}{(z-1)^3} dz, \\
 & \quad r) \int_{|z|=3} \frac{e^{zt}}{z^2+1} dz, \quad s) \int_{|z|=3} \frac{ze^{zt}}{(z+1)^3} dz.
 \end{aligned}$$

**6.2.** Compute the following integral in terms of  $r \in (0, 1) \cup (1, 2) \cup (2, \infty)$ :

$$I = \int_{|z|=r} \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z-1)(z-2)} dz.$$

**6.3.** Compute the following integral along the curve  $\gamma$  in the following cases:

$$\int_{\gamma} \frac{e^z}{z(1-z)^3} dz,$$

- a)  $\gamma$  is any curve with  $n(\gamma, 0) = 1, n(\gamma, 1) = 0$ .
- b)  $\gamma$  is any curve with  $n(\gamma, 0) = 0, n(\gamma, 1) = 1$ .
- c)  $\gamma$  is any curve with  $n(\gamma, 0) = 1, n(\gamma, 1) = 1$ .
- d)  $\gamma$  is any curve with  $n(\gamma, 0) = 0, n(\gamma, 1) = 0$ .

**6.4.** a) Prove the *Gauss' mean value theorem*: If  $f$  is holomorphic on a domain containing  $\{|z - z_0| \leq r\}$ , then

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + re^{it}) dt.$$

*Hint:* Use Cauchy's integral formula along the circumference with center  $z_0$  and radius  $r$ .

b) Prove that

$$\int_0^{2\pi} \cos(\cos \theta) \cosh(\sin \theta) d\theta = 2\pi.$$

*Hint:* Apply the previous item to the function  $f(z) = \cos z$ .

c) Deduce the mean value theorem for harmonic functions: If  $u$  is a harmonic function on a domain containing  $\{|z - z_0| \leq r\}$ , then

$$u(z_0) = \frac{1}{2\pi} \int_0^{2\pi} u(z_0 + re^{it}) dt.$$

*Hint:* Every harmonic function on a simply connected set  $D$  has a conjugate harmonic function on  $D$ .

**6.5.** a) Let  $f(z)$  be an holomorphic function on  $\{z \in \mathbf{C} : |z| < R_0\}$ . Prove that if  $|a| < R < R_0$ , and  $\gamma = \{z \in \mathbf{C} : |z| = R\}$ , then

$$f(a) = \frac{1}{2\pi i} \int_{\gamma} \frac{R^2 - |a|^2}{(z-a)(R^2 - z\bar{a})} f(z) dz.$$

*Hint:* Write the rational function on  $z$  as a sum of simple fractions and apply Cauchy's integral formula.

b) Deduce the *Poisson formula* for holomorphic functions (by using a)), for  $0 \leq r < R$  and  $0 \leq \theta < 2\pi$ :

$$f(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} \frac{R^2 - r^2}{R^2 - 2Rr \cos(\theta - \varphi) + r^2} f(Re^{i\varphi}) d\varphi.$$

*Hint:* Consider  $a = re^{i\theta}$  and  $z = Re^{i\varphi}$ .

c) Deduce the *Poisson formula* for harmonic functions  $u$  on  $\{z \in \mathbf{C} : |z| < R_0\}$  (by using b)) for  $0 \leq r < R$  and  $0 \leq \theta < 2\pi$ :

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} \frac{R^2 - r^2}{R^2 - 2Rr \cos(\theta - \varphi) + r^2} u(Re^{i\varphi}) d\varphi.$$

*Hint:* Every harmonic function on a simply connected set  $D$  has a conjugate harmonic function on  $D$ .

**6.6.** a) Assume that  $\gamma, \gamma_1, \gamma_2, \dots, \gamma_n$  are simple closed curves such that  $\gamma_1, \gamma_2, \dots, \gamma_n$  are contained in the interior of  $\gamma$ . If  $f$  is an holomorphic function on the closure of the region  $\Omega$  with  $\partial\Omega = \cup_{k=1}^n \gamma_k \cup \gamma$ , prove that:

$$\int_{\gamma} f(z) dz = \sum_{k=1}^n \int_{\gamma_k} f(z) dz.$$

*Hint:* Add some paths to the integral in order to write  $\Omega$  as a union of simply connected domains.

b) Under the hypotheses in the previous item, prove the following Cauchy's formula, for every  $a \in \Omega$ :

$$\int_{\gamma} \frac{f(z) dz}{z-a} - \sum_{k=1}^n \int_{\gamma_k} \frac{f(z) dz}{z-a} = 2\pi i f(a).$$

*Hint:* Use the previous item with the function  $(f(z) - f(a))/(z-a)$ .

c) Let  $f$  be an holomorphic function on  $\{z \in \mathbf{C} : 0 < |z| < R\}$ . Prove that the value of  $\int_0^{2\pi} f(re^{it}) dt$ , with  $0 < r < R$ , is independent of  $r$ . If  $f$  is holomorphic on the whole disk, compute the value of  $\int_0^{2\pi} f(re^{it}) dt$ .

*Hint:* Use item a).

**6.7.** Prove *Liouville's theorem*: If  $f$  is an entire function (i.e., holomorphic on  $\mathbf{C}$ ) satisfying  $|f(z)| \leq M$ , for every  $z \in \mathbf{C}$ , then  $f$  is constant.

*Hint:* Prove:

a)  $|f'(a)| \leq M/r^2$  for every  $a \in \mathbf{C}$ .

*Hint:* Use the Cauchy's inequality for  $f'$ .

b)  $f'(a) = 0$  for every  $a \in \mathbf{C}$ .

*Hint:* Take the limit in the inequality of item a) as  $r$  goes to some appropriate value.

**6.8.** Prove the *Fundamental Theorem of Algebra*: Every polynomial (with complex coefficients) of degree  $n \geq 1$  has  $n$  complex zeros (taking into account the multiplicity of its zeros).

*Hint:* It suffices to prove that the polynomial  $P$  has a zero. Seeking for a contradiction assume that  $P \neq 0$  and apply Liouville's theorem to the function  $1/P$ .

**6.9.** Let  $f$  be an entire function. Prove the following statements by using Liouville's theorem:

- If  $|f| \geq 1$ , then  $f$  is constant.
- If  $\operatorname{Re} f \geq 0$ , then  $f$  is constant.
- If  $\operatorname{Im} f \leq 1$ , then  $f$  is constant.
- If  $\operatorname{Re} f$  does not have zeros, then  $f$  is constant.
- If there exists a straight line that does not intersect the image of  $f$ , then  $f$  is constant.

**6.10.** Prove that the following functions are entire if they are defined in an appropriate way at their singular points:

$$\text{a) } \frac{\sin z}{z}, \quad \text{b) } \frac{e^z - 1 - z}{z^2}, \quad \text{c) } \frac{\sin(\pi z)}{z^3 - z}, \quad \text{d) } \frac{\sin(\pi z^2)}{\sin(\pi z)}.$$

**6.11.** Prove that  $F(z) = \int_0^1 e^{-z^2 x^2} dx$  is an entire function, and compute  $F'(z)$ .

**6.12.** If  $f : [0, \infty) \rightarrow \mathbf{C}$  is a function with  $\lim_{x \rightarrow \infty} f(x) e^{-ax} = 0$  for some  $a \in \mathbf{R}$  and  $f \in L^1([0, n])$  for every  $n$ , then the Laplace transform of  $f$  is defined as

$$Lf(z) = \int_0^\infty f(x) e^{-zx} dx.$$

Prove that  $f(x) e^{-zx}$  is integrable on  $[0, \infty)$  for each complex number  $z$  with  $\operatorname{Re} z > a$  and that  $Lf$  is holomorphic on the halfplane  $\{z \in \mathbf{C} : \operatorname{Re} z > a\}$ . Prove that the properties of the Laplace transform as a function of a real variable also hold for the Laplace transform as a function of a complex variable.

**6.13.** It can be proved that if  $f$  is a "good enough" function, then we can obtain  $f$  from its Laplace transform by the following Mellin's inverse formula:

$$f(x) = \lim_{T \rightarrow \infty} \frac{1}{2\pi i} \int_{\gamma_{t,T}} e^{zx} (Lf)(z) dz,$$

where if  $\lim_{s \rightarrow \infty} f(s) e^{-as} = 0$ , then  $\gamma_{t,T}$  is the vertical segment  $\gamma_{t,T} = \{z \in \mathbf{C} : \operatorname{Re} z = t, \operatorname{Im} z \in [-T, T]\}$  oriented starting at  $t - iT$  and ending at  $t + iT$ , with  $t > a$ . Alternatively, we can choose  $t$  greater than the real part of all singularities of  $F(z)$ .

Compute  $f(x)$  if:

$$(Lf)(z) = \frac{1}{z-3}, \quad (Lf)(z) = \frac{z}{(z-1)^2(z^2+3z-10)}, \quad (Lf)(z) = \frac{1}{z^2(z^2+2z+2)},$$

and check that its value is independent on the choice of  $t > a$ .

*Hint:* Apply Cauchy's integral formula for an appropriate closed curve containing  $\gamma_{t,T}$  and a part of circumference  $C_T$  joining the endpoints of  $\gamma_{t,T}$ . Prove that the integral along  $C_T$  goes to 0 as  $T$  goes to  $\infty$ .

**6.14.** Let  $f$  be an holomorphic function on a simply connected domain  $D$ , such that  $f(z) \neq 0$  for every  $z \in D$ .

a) Prove that for any  $z_0 \in D$ , the function  $L(z) = \int_{z_0}^z f'(w)/f(w) dw$  is well defined (i.e., the value of  $L(z)$  is independent of the curve joining  $z_0$  with  $z$ ) and so, it is possible to define the function  $\log f(z)$  in such a way that it is holomorphic on  $D$ .

b) Prove that for any  $\alpha \in \mathbf{C}$ , it is possible to define the function  $f(z)^\alpha$  in such a way that it is holomorphic on  $D$ .

c) Do items a) or b) hold in general if  $D$  is not simply connected?

d) Find a domain  $D$  such that  $(1 - z^2)^{-1/2}$  can be defined as a holomorphic function on  $D$ . Is  $\arcsin z$  holomorphic on that domain?

e) Find a domain  $D$  such that  $\arctan z$  can be defined as a holomorphic function on  $D$ .

Hints: a) Use that  $\int_{\gamma}^z f'(w)/f(w) dw = 0$  for any closed curve  $\gamma$  in  $D$ . b) Use the previous item.

**6.15.** Let  $f$  be a meromorphic function on a domain  $D$  (i.e., a holomorphic function on  $D$  except for a set of isolated points, which are poles of  $f$ ).

a) Prove that if  $f$  has a zero of order  $k$  at  $a$ , then the function

$$\frac{f'(z)}{f(z)} - \frac{k}{z-a}$$

is holomorphic on a neighborhood of  $a$ .

b) Prove that if  $f$  has a pole of order  $k$  at  $a$ , the function

$$\frac{f'(z)}{f(z)} + \frac{k}{z-a}$$

is holomorphic on a neighborhood of  $a$ .

Hints: a) If  $f$  has a zero of order  $k$  at  $a$ , then  $f(z) = (z-a)^k g(z)$  where  $g$  is a holomorphic function on a neighborhood of  $a$  with  $g(a) \neq 0$ . b) If  $f$  has a pole of order  $k$  at  $a$ , then  $f(z) = g(z)/(z-a)^k$  where  $g$  is a holomorphic function on a neighborhood of  $a$  with  $g(a) \neq 0$ .

**6.16.** a) Let  $f$  be a meromorphic function on a simply connected domain  $D$  with zeros  $a_1, a_2, \dots, a_r$  and poles  $b_1, b_2, \dots, b_s$  (in each list appear the zeros and the poles taking into account their multiplicities, i.e., if a zero or a pole has order  $k$ , it appears  $k$  times in the list). Prove that if  $\gamma$  is a closed curve contained on  $D$  with  $\gamma \cap \{a_1, a_2, \dots, a_r, b_1, b_2, \dots, b_s\} = \emptyset$ , and  $\Gamma = f \circ \gamma$  is the image of  $\gamma$  by  $f$ , then

$$\frac{1}{2\pi i} \int_{\Gamma} \frac{dw}{w} = \frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = \sum_{j=1}^r n(\gamma, a_j) - \sum_{k=1}^s n(\gamma, b_k).$$

This result is known as the *argument principle*. In particular, if  $\gamma$  is a Jordan curve in  $D$  enclosing a simply connected domain  $D_{\gamma} \subset D$ , the argument principle gives that the above integral is equal to the number of zeros of  $f$  on  $D_{\gamma}$  minus the number of poles of  $f$  on  $D_{\gamma}$  (taking into account their multiplicities).

Hint: In order to prove the first equality, note that if  $\gamma(t)$  is a parametrization of  $\gamma$ , then  $f(\gamma(t))$  is a parametrization of  $\Gamma$ . In order to prove the second equality, use the previous exercise to show that the following function is holomorphic on the domain  $D$

$$\frac{f'(z)}{f(z)} - \sum_{j=1}^r \frac{1}{z-a_j} + \sum_{k=1}^s \frac{1}{z-b_k}.$$

b) Use the previous item to compute

$$i) \int_{|z|=2} \tan z dz, \quad ii) \int_{|z|=2} \frac{dz}{\sin z \cos z}.$$

Hints: i) Consider the function  $f(z) = \cos z$ . ii) Consider the function  $f(z) = \operatorname{cosec} 2z + \cotan 2z$ .

Solutions: i)  $2\pi i$ , ii)  $-2\pi i$ .

**6.17.** Let  $f, g$  be two holomorphic functions on a domain  $D$ , and  $\gamma$  a Jordan curve (i.e., a simple closed curve) in  $D$  surrounding a simply connected domain  $D_{\gamma} \subset D$ . If  $f, g$  satisfy the inequality  $|f(z) - g(z)| < |f(z)|$  for every  $z \in \gamma$ , prove that:

a) The function  $F = g/f$  does not have zeros nor poles in the curve  $\gamma$ , i.e.,  $f$  and  $g$  do not have zeros in  $\gamma$ .

b) If  $\Gamma$  is the image by  $F$  of  $\gamma$ , then  $\int_{\Gamma} dw/w = 0$ .

Hint: Since  $|g(z)/f(z) - 1| < 1$  on  $\gamma$ , we have  $|w - 1| < 1$  on  $\Gamma$ .

c) Prove that  $f(z)$  and  $g(z)$  have the same number of zeros on  $D_\gamma$  (taking into account their multiplicities). This result is known as *Rouché's theorem*.

*Hint:* Use the argument principle.

**6.18.** Apply Rouché's theorem in order to solve the following problems:

a) How many roots does the equation  $z^7 - 2z^5 + 6z^3 - z + 1 = 0$  have in the unit disk  $\mathbf{D} = \{|z| < 1\}$ ?

*Hint:* Consider  $g(z) = z^7 - 2z^5 + 6z^3 - z + 1$  and choose  $f(z)$  as the monomial of  $g(z)$  with greatest modulus on  $\{|z| = 1\}$ .

b) How many roots does the equation  $z^7 - 2z^5 + 6z^3 - z + 1 = 0$  have in the disk  $\{|z| < 2\}$ ?

c) How many roots does the equations  $z^9 - 2z^6 + z^2 - 8z - 2 = 0$ ,  $2z^5 - z^3 + 3z^2 - z + 8 = 0$ ,  $z^7 - 5z^4 + z^2 - 2 = 0$ , have in  $\mathbf{D}$ ?

d) How many roots does the equation  $z^4 - 6z + 3 = 0$  have in the disk  $\{|z| < 2\}$ ? And in  $\mathbf{D}$ ? And in the annulus  $\{1 < |z| < 2\}$ ?

e) How many roots does the equation  $z^4 - 5z + 1 = 0$  have in  $\mathbf{D}$ ? And in the annulus  $\{1 < |z| < 2\}$ ?

f) How many roots does the equation  $z^4 - 8z + 10 = 0$  have in  $\mathbf{D}$ ? And in the annulus  $\{1 < |z| < 3\}$ ?

g) How many roots does the equation  $z^n + az^2 + bz + c = 0$  have in  $\mathbf{D}$ , if  $|a| > |b| + |c| + 1$  and  $n \in \mathbf{N}$ ?

h) How many roots does the equation  $z = f(z)$  have in  $\mathbf{D}$ , if  $f$  is an holomorphic function satisfying  $|f(z)| < 1$  if  $|z| \leq 1$ ?

i) How many roots does the equation  $e^z - 4z^n + 1 = 0$  have in  $\mathbf{D}$ , if  $n \in \mathbf{N}$ ?

**6.19.** Let  $f$  be an holomorphic function on  $\mathbf{D}$  satisfying  $|f(z)| < 1$  for every  $z \in \mathbf{D}$  and  $f(0) = 0$ . Prove that:

a) The function  $g(z) = f(z)/z$  is holomorphic on  $\mathbf{D}$ .

b) For each  $0 < r < 1$  we have  $|g(z)| \leq 1/r$  if  $|z| \leq r$ .

c)  $|f(z)| \leq |z|$  for every  $z \in \mathbf{D}$  and  $|f'(0)| \leq 1$ .

d) If there exists a point  $z_0 \in \mathbf{D}$  such that  $|f(z_0)| = |z_0|$  (or  $|f'(0)| = 1$ ), then  $f(z) = cz$  where  $c$  is a complex number with  $|c| = 1$ .

These results are known as *Schwarz Lemma*.

**6.20.** Prove the *minimum modulus principle*: If  $f(z)$  is an holomorphic function on the domain  $D$ ,  $f(z) \neq 0$  for every  $z \in D$ , and  $|f(z)|$  attains its minimum value at a point in  $D$ , then  $f(z)$  is constant.

**6.21.** Study if there exists an holomorphic function on  $\mathbf{D}$  such that on the points  $1/n$  ( $n = 1, 2, 3, \dots$ ) take the values:

a)  $0, 1, 0, 1, 0, 1, \dots$

b)  $0, 1/2, 0, 1/4, 0, 1/6, \dots, 0, 1/(2k), \dots$

c)  $1/2, 1/2, 1/4, 1/4, 1/6, 1/6, \dots, 1/(2k), 1/(2k), \dots$

d)  $1/2, 2/3, 3/4, 4/5, 5/6, 6/7, \dots, n/(n+1), \dots$